Healthy Kentuckians 2010: Schools and Tobacco Use

Healthy Kentuckians 2010 recommends increasing the proportion of schools with tobacco-free environments and increasing the proportion of schools that provide research-based tobacco use prevention curricula. Research shows that the school health programs that use skills training approaches are the most effective in preventing tobacco use. Programs and policies that prevent the onset of smoking during middle school years are a crucial part of a comprehensive tobacco prevention program. In addition, programs targeting high school students reinforce the earlier messages and can provide an opportunity for youth to develop leadership skills within their communities.

Schools and Tobacco Use

Smoking on School Property

- ✓ Overall, 6% of middle school students and 19% of high school students report smoking cigarettes on school property during the past 30 days.
- ✓ High school smokers are more likely to smoke cigarettes on school property than middle school smokers.




Figure 43. Students who smoked cigarettes on school property during the past 30 days

Background and Significance

Nationally, 2% of middle school students and 9% of high school students smoked cigarettes on school property in the past 30 days. A higher proportion of Kentucky youth smoke on school property than U.S. youth. Schools that have strong, consistently enforced policies on tobacco use can help students decide not to use tobacco. A tobacco-free environment has many benefits including decreased fires and discipline problems related to smoking and easier maintenance of school facilities and grounds. Schools have an important opportunity to promote tobacco-free norms by developing and enforcing tobacco-free policies including all athletic facilities, vehicles, and school grounds.

Schools and Tobacco Use

Practicing Ways to Say "No" to Tobacco

- ✓ Overall, 43% of middle school students and 15% of high school students practice ways to say "NO" to tobacco at school.
- ✓ Middle school students are more likely to practice ways to say "NO" to tobacco use at school than high school students.




Figure 44. Students who practiced ways to say "NO" to tobacco by gender

Background and Significance

The findings reported here are consistent with the fact that about one in four middle and high schools in Kentucky report using at least one research-based alcohol, tobacco, and drug curriculum, and middle schools are more likely to teach tobacco prevention than high schools. School health programs that use skills training approaches have been shown to be effective in reducing the onset of tobacco use. The CDC recommends that schools provide tobacco use prevention education in kindergarten through 12th grade and that it be especially intensive in middle school and reinforced in high school.